

## Product Information: Jevity® 1.5 Cal

For more information, contact your Abbott Nutrition Representative or visit  
[www.abbottnutrition.com](http://www.abbottnutrition.com)

# Jevity® 1.5 Cal

Complete, Balanced Nutrition® with fiber and NutraFlora® scFOS®

- 1 JEVITY 1.5 CAL is a calorically dense formula with a unique fiber blend that provides Complete, Balanced Nutrition®.
- 1 For tube feeding.
- 1 For supplemental or sole-source nutrition.
- 1 May be used for oral feeding of patients with altered taste perception.
- 1 Use under medical supervision.
- 1 Concentrated calories (1.5 Cal/mL) to help tube-fed patients gain and maintain healthy weight.
- 1 22 grams of dietary fiber in 1 Liter (12 g of a unique soluble and insoluble fiber blend and 10 g of short-chain fructooligosaccharides [scFOS]), which helps meet daily fiber needs. Fiber helps moderate bowel function.
- 1 Includes 15 g NutraFlora® scFOS® /1500 mL. scFOS are prebiotics that stimulate the growth of beneficial bacteria in the colon.
- 1 Meets or exceeds 100% of the 1995 RDIs for 24 essential vitamins and minerals in 1500 Cal (1 Liter).
- 1 Halal.
- 1 Kosher.
- 1 Gluten-free.
- 1 Suitable for lactose intolerance.



## Safety Precautions

- 1 Not for people with galactosemia.
- 1 Not for IV use.

\* NutraFlora® scFOS® is not a registered trademark of Abbott Laboratories.

## Ingredients

### Liquid Unflavored:

Water, Corn Maltodextrin, Corn Syrup Solids, Sodium & Calcium Caseinates, Canola Oil, Corn Oil, Soy Protein Isolate, Short-Chain Fructooligosaccharides, Medium-Chain Triglycerides, Oat Fiber, Soy Fiber, Potassium Citrate, Calcium Phosphate, Gum Arabic, Sodium Citrate, Magnesium Phosphate, Soy Lecithin, Magnesium Chloride, Cellulose Gum, Ascorbic Acid, Choline Chloride, Potassium Chloride, Taurine, L-Carnitine, Zinc Sulfate, dl-Alpha-Tocopheryl Acetate, Ferrous Sulfate, Niacinamide, Calcium Pantothenate, Manganese Sulfate, Copper Sulfate, Thiamine Hydrochloride, Pyridoxine Hydrochloride, Riboflavin, Vitamin A Palmitate, Folic Acid, Biotin, Chromium Chloride, Sodium Molybdate, Potassium Iodide, Sodium Selenate, Phylloquinone, Vitamin B12, and Vitamin D3.

**Allergens:** Contains milk and soy ingredients.

## Availability

List Number	Item
57333	Jevity 1.5 Cal / 8-fl-oz (237-mL) Can / 24 ct
62681	Jevity 1.5 Cal / 1000-mL (1.1-QT) Ready-To-Hang Prefilled Container / 8 ct
62689	Jevity 1.5 Cal / 1500-mL (1.6-QT) Ready-To-Hang Prefilled Container / 6 ct
64628	Jevity 1.5 Cal / 8-fl-oz (237-mL) Carton / 24 ct

For more information, contact your Abbott Nutrition Representative or visit  
[www.abbottnutrition.com](http://www.abbottnutrition.com)

## Nutrition Information - Liquid Unflavored

	8 fl oz (237 mL)		1000 mL (1.1 QT)		1500 mL (1.6 QT)	
	Value	%RDI / %DV	Value	%RDI / %DV	Value	%RDI / %DV
Protein, g	15.1		63.8		95.7	
Fat, g	11.8		49.8		74.7	
Carbohydrate, g	51.1*		215.7		323.6	
Dietary Fiber, g	5.3*		22.0*		33.0*	
L-Carnitine, mg	36		150		225	
Taurine, mg	36		150		225	
Water, g/mL/cc	180		760		1140	
Calories	355		1500		2250	
Vitamin A, IU	1185		5000		7500	
Vitamin D, IU	95		400		600	
Vitamin E, IU	11		45		68	
Vitamin K, mcg	19		80		120	
Vitamin C, mg	72		300		450	
Folic Acid, mcg	145		600		900	
Thiamin (Vitamin B1), mg	0.54		2.3		3.4	
Riboflavin (Vitamin B2), mg	0.61		2.6		3.9	
Vitamin B6, mg	0.72		3.0		4.5	
Vitamin B12, mcg	2.2		9.0		14	
Niacin, mg	7.2		30		45	
Choline, mg	145		600		900	
Biotin, mcg	110		450		675	
Pantothenic Acid, mg	3.6		15		23	
Sodium, mg	330		1400		2100	
Sodium, mEq	14.3		60.9		91.3	
Potassium, mg	510		2150		3225	
Potassium, mEq	13.1		55.1		82.7	
Chloride, mg	320		1360		2040	
Chloride, mEq	9.0		38.4		57.6	
Calcium, mg	285		1200		1800	
Phosphorus, mg	285		1200		1800	
Magnesium, mg	95		400		600	
Iodine, mcg	36		150		225	
Manganese, mg	1.2		5.0		7.5	
Copper, mg	0.48		2.0		3.0	
Zinc, mg	5.4		23		34	
Iron, mg	4.3		18		27	
Selenium, mcg	17		70		105	
Chromium, mcg	29		120		180	
Molybdenum, mcg	36		150		225	

For more information, contact your Abbott Nutrition Representative or visit  
[www.abbottnutrition.com](http://www.abbottnutrition.com)

## Liquid Unflavored Footnotes & References

### Per 8 fl oz (237 mL)

\*2.9 g of total dietary fiber from a unique fiber blend and 2.4 g of short-chain fructooligosaccharides

### Per 1000 mL (1.1 QT)

\*12.0 g of total dietary fiber from a unique fiber blend and 10.0 g of fructooligosaccharides.

### Per 1500 mL (1.6 QT)

\*18.0 g of total dietary fiber from a unique fiber blend and 15.0 g of fructooligosaccharides.

## Preparation

### Instructions for Use:

#### 8-fl-oz container

- 1 Store unopened at room temperature; avoid extreme temperatures.
- 1 Shake well prior to opening.
- 1 Clean top of can thoroughly before opening.
- 1 Once opened, reclose or cover, refrigerate and **use within 48 hours**.

#### Oral Feeding:

- 1 May be used for total or supplemental nutrition.
- 1 May be fed at room temperature or chilled.

#### Tube Feeding:

- 1 Follow physician's instructions.
- 1 Adjust flow rate and volume according to patient's condition and tolerance.
- 1 Feed at room temperature using a feeding pump or syringe.
- 1 Additional fluid requirements should be met by giving water between or after feedings or when flushing the tube.
- 1 Avoid contamination during preparation and use.

### Ready-To-Hang Container

All liquid medical foods, regardless of type of administration system, require careful handling because they can support microbial growth. Follow these instructions for clean technique and proper setup to reduce the potential for microbial contamination.

NOTE: Failure to follow the Instructions for Use increases the potential for microbial contamination and reduces hangtime.

- 1 Administer product at room temperature.
- 1 THOROUGHLY wash hands with soap and water before handling container or feeding set.
- 1 Turn container upside down and SHAKE VIGOROUSLY, using a twisting motion for at least 10 seconds.
- 1 DO NOT touch any part of the container or feeding set that comes into contact with the formula.
- 1 When initiating feeding, follow physician's instructions. Adjust flow rate and volume according to patient's condition and tolerance.
- 1 Additional fluid requirements should be met by giving water between or after feedings or when flushing the tube.

#### For Use with Enteral Feeding Pumps:

- 1 Remove dust cover from RTH Safety Screw Cap.
- 1 Remove dust cover from feeding set connector.
- 1 Insert feeding set connector into port of RTH Safety Screw Cap and completely pierce foil.
- 1 Turn connector collar clockwise until it is securely fastened.
- 1 Close clamp on set before inverting container.
- 1 Invert container and suspend, using hanging ring on bottom of container.
- 1 Follow directions for use provided by manufacturer of feeding sets.
- 1 Unless a shorter hang time is specified by the set manufacturer, hang product for up to 48 hours after initial connection when clean technique and only one new set are used. Otherwise hang for no more than 24 hours.