



Banatrol[®] plus

With Bimuno[®] Prebiotic

For Diarrhea and Loose Stools A Medical Food

Formulated for patients under medical supervision experiencing diarrhea and loose stools associated with the flu, antibiotics, tube feeding, oncology treatment and Clostridium difficile (*C. diff.*) Does not affect gut motility. Can be used as soon as *C. diff.* is suspected.

Banana
Natural Flavor

Net Weight 0.38oz. (10.75g)

Item 18470

DOSING AND ADMINISTRATION: Oral: Diarrhea: One packet of Banatrol[®] Plus TID (three times daily) added to a moist food such as applesauce or yogurt or to a beverage. Titrate the dose as needed with a minimum of one packet/day and maximum of six packets/day. Continue until diarrhea is resolved. **Loose Stools:** One packet once or twice a day as needed. **Tube Feeding-G-Tube or J-tube:** One packet of Banatrol[®] Plus TID every six to eight hours of feeding. Add 120 mL of room-temperature water to a 240-mL cup. Slowly stir in the contents of one packet of Banatrol Plus. Stir until dissolved. Administer slowly via syringe down tube. Depending on the size of the feeding tube, you may need to use the syringe plunger to gently push mixture down the tube. Flush with 30 mL water before and after administration. Do not mix with tube feeding formula.

Nutrition Facts

Serving Size: 1 Packet (10.75 g)

Amount Per Serving

Calories 40 Calories from Fat 0

% Daily Values*

Total Fat 0g **0%**

Saturated Fat 0g **0%**

Trans Fat 0g

Cholesterol 0mg **0%**

Sodium 3mg **0%**

Potassium 120mg **3%**

Phosphorus 5mg **0%**

Total Carbohydrate 9g **3%**

Dietary Fiber 2g **8%**

Sugars 5g

Protein <1g

Vitamin A 0% • Vitamin C 0%

Calcium 0% • Iron 0%

* Percent Daily Values are based on a 2,000 calorie diet.

When administering Banatrol Plus via feeding tube, we recommend a size-14 French feeding tube or larger to reduce the risk of clogging the tube.

For Smaller French-size tubes and additional instructions see our website:
www.medtrition.com/products/banatrol-plus/dosing
Can be used for long term maintenance if needed.

Does not cause constipation.

Contraindications: Latex-fruit syndrome

INGREDIENTS: Banana Flakes, Bimuno[®] (transgalactooligosaccharides, lactose, glucose, gum acacia, galactose). **CONTAINS MILK.**

Contains <0.550 mcg of Vitamin K per serving.

Low in Lactose: 0.72 g/serving

Osmolality: 240 mOsm/kg water

HCPCS: B4104, Billing Code: 94688-0184-70

Not a sole source of nutrition.

A Medical Food. Use only under care of a physician.

NOT FOR PARENTERAL USE

Distributed by
Medtrition, Inc.

P.O. Box 5387

Lancaster, PA 17606

Toll Free 877.271.3570

info@medtrition.com

Product of USA



For clinical studies which support the use of this product as directed, visit www.medtrition.com
Bimuno is a registered trademark of Clasado, Inc. ©2016 Medtrition, Inc. RH6H