



NDC 57896-226-01
THE SENSIBLE CHOICE
FOR PRICE AND HEALTH

**Extra
Strength**

Pain-Reliever

Acetaminophen 250 mg
Aspirin 250 mg (NSAID)
Caffeine 65 mg

COMPARE TO THE ACTIVE
INGREDIENTS IN EXCEDRIN®

100 Caplets

Tamper Evident: Do not use if imprinted seal under cap is missing or broken.

Drug Facts

Active Ingredients (in each caplet)

Acetaminophen 250 mg	Pain reliever
Aspirin 250 mg (NSAID)*	Pain reliever
Caffeine 65 mg	*Nonsteroidal anti-inflammatory drug

Uses ■ Temporarily relieves minor aches and pains

Warnings

Reye's Syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Aspirin may cause a severe allergic reaction which may include: ■ hives ■ facial swelling ■ asthma (wheezing) ■ shock

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take ■ more than directed ■ with other drugs containing acetaminophen ■ 3 or more alcoholic drinks every day while using this product

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you ■ are age 60 or older ■ have had stomach ulcers or bleeding problems ■ take a blood thinning (anticoagulant) or steroid drug ■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) ■ have 3 or more alcoholic drinks every day while using this product ■ take more or for a longer time than directed

Caffeine warning: The recommended dose of this product contains about as much caffeine as a cup of coffee. Limit the use of caffeine-containing medications, foods, or beverages while taking this product because too much caffeine may cause nervousness, irritability, sleeplessness, and, occasionally, rapid heartbeat.

*This product is not manufactured or distributed by the owner of the registered trademark Excedrin®.

Distributed By: Geni-Care Pharmaceuticals Corp.
1650 63rd Street, Brooklyn, NY 11204



3 57896 22601 5
PEEL HERE FOR MORE DRUG FACTS

Drug Facts (continued)

Do not use ■ If you have ever had an allergic reaction to aspirin or any other pain reliever/fever reducer ■ with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

Ask a doctor before use if ■ you have liver disease ■ stomach bleeding warning applies to you ■ you have a history of stomach problems, such as heartburn ■ you have high blood pressure, heart disease, liver cirrhosis, or kidney disease ■ you are taking a diuretic ■ you have asthma

Ask a doctor or pharmacist before use if you are taking ■ any other drug containing an NSAID (prescription or nonprescription) ■ blood thinning (anticoagulant) or steroid drug ■ a prescription drug for diabetes, gout, or arthritis ■ any other drug, or are under a doctor's care for any chronic condition

Stop use and ask a doctor if ■ an allergic reaction occurs. Seek medical help right away.

■ you experience any of the following signs of stomach bleeding: ☐ faint/flat ☐ vomit blood ☐ have bloody or black stools

■ have stomach pain that does not get better ■ ringing in the ears or loss of hearing occurs ■ painful area is red or swollen ■ pain gets worse or lasts for more than 10 days ■ fever gets worse or lasts for more than 3 days ■ any new symptoms appear

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children.

Overdose Warning: Taking more than the recommended dose (or doses) may cause liver damage. In case of overdose get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions ■ do not use more than directed (see overdose warning)

- drink a full glass of water with each dose
- adults and children 12 years and over: take 2 caplets every 6 hours; not more than 8 caplets in 24 hours
- children under 12: ask a doctor

Other information

■ store at room temperature 15° - 25°C (59° - 77°F)

Inactive ingredients camellid wax, cellulose, hypromellose, mineral oil, povidone, silica, sodium starch glycolate, starch, stearic acid, titanium dioxide.

STOP PEELING