Pulmocare®

Therapeutic nutrition for people With COPD

PULMOCARE is designed for people with chronic obstructive pulmonary disease (COPD), cystic fibrosis, or respiratory failure who may benefit from a high-calorie, modified carbohydrate and fat enteral formula that may help reduce diet-induced carbon dioxide production.[1,2,3,4] For tube or oral feeding. For supplemental or sole-source nutrition. Use under medical supervision.

[1] Angelillo VA, et al. Ann Intern Med. 1985;103:883-885.
[2] Goldstein SA, et al. Clin Chest Med. 1986;7(1):141-151.
[3] Cai B, et al. Nutrition. 2003; 19(3):229-32.
[4] Sklarek H, et al. Am Rev Respir Dis. 1987;135-A359.



Features

- Provides complete and balanced nutrition.
- 20% of fat as MCTs to enhance fat absorption.
- Elevated antioxidants vitamins C & E, and beta-carotene.
- Kosher.
- Halal.
- Gluten-free.
- Suitable for lactose intolerance.

Safety Precautions

- Not for IV use.
- Not suitable for people with galactosemia.

Availability

List Number	Item
62725	Pulmocare / 33.8 fl oz (1 L) Ready-To-Hang Prefilled Container / 8 ct

Ingredients

Ready to Hang Liquid

Water, Sodium Caseinate, Sugar, Canola Oil, Corn Maltodextrin, Medium Chain Triglycerides, Corn Oil, Calcium Caseinate, High Oleic Safflower Oil. Less than 0.5% of: Natural & Artificial Flavor, Calcium Phosphate, Magnesium Phosphate, Soy Lecithin, Potassium Citrate, Potassium Chloride, Sodium Citrate, Salt, Choline Chloride, Ascorbic Acid, L-Carnitine, Taurine, Gellan Gum, Ferrous Sulfate, dl-Alpha-Tocopheryl Acetate, Zinc Sulfate, Niacinamide, Calcium Pantothenate, Manganese Sulfate, Copper Sulfate, Pyridoxine Hydrochloride, Thiamine Hydrochloride, Riboflavin, Vitamin A Palmitate, Beta-Carotene, Folic Acid, Chromium Chloride, Sodium Molybdate, Potassium Iodide, Sodium Selenate, Phylloquinone, Biotin, Vitamin D3, and Vitamin B12.

Contains milk and soy ingredients.

Nutrition Information

	33.8 fl oz (1 L)	
	Value	%DV
Protein, g	62.6	
Fat, g	93.3	
Carbohydrate, g	105.7	
Total Sugars, g	54	
Added Sugars, g	54	

Abbott Nutrition Abbott Laboratories
Columbus, OH 43219

1-800-227-5767

For more information contact your Abbott Nutrition Representative or visit:

www.abbottnutrition.com

© 2022 Abbott Laboratories Inc.

Last updated: 10/07/2023

Added Sugars, g	54	
L-Carnitine, mg	160	
Taurine, mg	160	
Water, mL	785	
Calories	1500	
Vitamin A, mcg RAE	930*	
Vitamin D, mcg	20.3	
Vitamin E, mg	27	
Vitamin K, mcg	122	
Vitamin C, mg	321	
Folate, mcg DFE	400	
Folic Acid, mcg	241	
Thiamin (Vitamin B1), mg	1.81	
Riboflavin (Vitamin B2), mg	1.90	
Vitamin B6, mg	2.19	
Vitamin B12, mcg	2.41	
Niacin, mg	16	
Choline, mg	550	
Biotin, mcg	35	
Pantothenic Acid, mg	10.1	
Sodium, mg	1520	
Potassium, mg	2320	
Chloride, mg	2320	
Calcium, mg	1310	
Phosphorus, mg	1260	
Magnesium, mg	420	
Iodine, mcg	152	
Manganese, mg	2.32	
Copper, mg	1.39	
Zinc, mg	11	
Iron, mg	18.1	
Selenium, mcg	55	
Chromium, mcg	63	
Molybdenum, mcg	72	

*320 mcg RAE of Vitamin A activity supplied by 0.8 mg Beta-Carotene.

Preparation

Instructions for Use:

8 fl oz Container

- Store unopened at room temperature; avoid extreme temperatures.
- Shake well prior to opening.

© 2022 Abbott Laboratories Inc.

- After opening, refrigerate and consume product within 48 hours. If consuming product directly from the container, refrigerate and consume within 24 hours.

Oral Feeding

- May be used for total or supplemental nutrition.
- May be fed at room temperature or chilled.

Tube Feeding

- Follow physician's instructions.
- Adjust flow rate and volume according to patient's condition and tolerance.
- Feed at room temperature using a feeding pump or syringe.
- Additional fluid requirements should be met by giving water between or after feedings or when flushing the tube.
- Avoid contamination during preparation and use.

Ready-To-Hang Container

All medical foods, regardless of type of administration system, require careful handling because they can support microbial growth. Follow these instructions for clean technique and proper setup to reduce the potential for microbial contamination.

NOTE: Failure to follow the **INSTRUCTIONS FOR USE** increases the potential for microbial contamination and may reduce hangtime.

- Administer product at room temperature.
- THOROUGHLY wash hands with soap and water before handling container or feeding set.
- Turn container upside down and SHAKE VIGOROUSLY, using a twisting motion for at least 10 seconds.
- DO NOT touch any part of the container or feeding set that comes into contact with the formula.
- When initiating feeding, follow physician's instructions. Adjust flow rate and volume according to patient's condition and tolerance.
- Additional fluid requirements should be met by giving water between or after feedings or when flushing the tube.

For Use with Enteral Feeding Pumps:

- Remove dust cover from RTH Safety Screw Cap.
- Remove dust cover from feeding set connector.
- Insert feeding set connector into port of RTH Safety Screw Cap and completely pierce foil.
- Turn connector collar clockwise until it is securely fastened.
- Close clamp on set before inverting container.
- Invert container and suspend, using hanging ring on bottom of container.

Precautions

- Follow directions for use provided by manufacturer of feeding sets.
- Unless a shorter hang time is specified by the set manufacturer, hang product for up to 48 hours after initial connection when clean technique and only one new set are used. Otherwise hang for no more than 24 hours.

© 2022 Abbott Laboratories Inc.