

PediaSure® 1.5 Cal with Fiber

Complete, Balanced Nutrition®

PEDIASURE 1.5 CAL WITH FIBER is a higher^{*} caloric density product designed to meet the higher energy requirements of pediatric patients who are at risk for malnutrition, require a higher caloric density, or have fluid restrictions. It provides a source of complete, balanced nutrition for children 1 to 13 years of age. For oral or tube feeding. May be used as a sole source of nutrition or as a supplement. Use under medical supervision.

^{*} Caloric value is 1.5 Cal/mL compared to PediaSure Enteral Formula 1.0 Cal at 1.0 Cal/mL.



Features

- Meets or exceeds 100% of the DRIs for protein and 25 essential vitamins and minerals
 - in 1000 mL for children 1 to 8 years of age.
 - in 1500 mL for children 9 to 13 years of age.
- Excellent source of high-quality protein designed to meet the acceptable macronutrient distribution range of protein for children 1 to 13 years of age.
- Vitamin D, calcium and phosphorus for strong bones.
- Prebiotics designed to help support digestive tract health^{1,2,3} and antioxidants (vitamins C & E and selenium) to support the immune system.
- DHA (32 mg per 8 fl oz) and lutein to support the brain and eyes.^{4,5}
- No artificial growth hormones.[†]
- Kosher.
- Halal.
- Gluten-free.
- Suitable for lactose intolerance.

¹ Bornet FR, et al. Nutr Rev. 2002;60(11):326-334.

[†] No significant difference has been shown between milk derived from rbST-treated and non-rbST-treated cows.

² Tokunaga T, et al. Bifidus. 1993;6(2):143-150.

³ Hidaka H, et al. Bifidobact Microflora. 1986;5:37-50.

⁴ Lauritzen L, et al. Prog Lipid Res. 2001;40:1-94.

⁵ Krinsky NI, et al. Annu Rev Nutr. 2003;23:171-201.

Safety Precautions

- Not intended for infants under 1 year of age unless specified by a physician.
- Not for IV use.
- Not for children with galactosemia.

Availability

List Number	Item
67376	PediaSure 1.5 Cal w Fiber / 33.8 fl oz (1 L) Ready-To-Hang Prefilled Container / 8 ct
67374	PediaSure 1.5 Cal w Fiber Vanilla / 8 fl oz (237 mL) Can / 24 ct

Ingredients

Vanilla Liquid

Water, Corn Maltodextrin, Blend of Vegetable Oils (High Oleic Safflower, Soy, Medium Chain Triglycerides), Milk Protein Concentrate, Soy Protein Isolate, Short-chain Fructooligosaccharides. Less than 0.5% of: Vitamins & Minerals (Potassium Citrate, Potassium Chloride, Magnesium Phosphate, Calcium Phosphate, Ascorbic Acid, Choline Chloride, Calcium Carbonate, Potassium Phosphate, Ferrous Sulfate, dl-Alpha-Tocopheryl Acetate, Zinc Sulfate, Niacinamide, Calcium Pantothenate, Manganese Sulfate, Thiamine Hydrochloride, Pyridoxine Hydrochloride, Riboflavin, Vitamin A Palmitate, Copper Sulfate, Folic Acid, Chromium Chloride, Potassium Iodide, Sodium Molybdate, Sodium Selenate, Biotin, Phyloquinone, Vitamin D3, Vitamin B12), Oat Fiber, Natural & Artificial Flavor, Soy Fiber, Cellulose Gel, Soy Lecithin, Tuna Oil*, Monoglycerides, Cellulose Gum, Potassium Hydroxide, Carrageenan, Inositol, Taurine, Sucralose, Salt, L-Carnitine, and Lutein.

* Source of DHA.


Contains milk and soy ingredients.

Nutrition Information

	8 fl oz (237 mL)	
	Value	%DV
Protein, g	14	
Fat, g	16	
Saturated Fat, g	3.5	
Trans Fat, g	0	
Polyunsaturated Fat, g	5	
Monounsaturated Fat, g	7	
Cholesterol, mg	10	
Carbohydrate, g	39	
Dietary Fiber, g	3	
scFOS, g	1.7	
Total Sugars, g	3	
Added Sugars, g	3	
L-Carnitine, mg	4.0	
Taurine, mg	18	
Water, mL	185	
Calories	350	
Vitamin A, mcg RAE	140	
Vitamin D, mcg	6.0	
Vitamin E, mg	3.0	
Vitamin K, mcg	18	
Vitamin C, mg	23	
Folate, mcg DFE	60	
Folic Acid, mcg	36	
Thiamin (Vitamin B1), mg	0.3	
Riboflavin (Vitamin B2), mg	0.33	
Vitamin B6, mg	0.34	
Vitamin B12, mcg	0.48	
Niacin, mg NE	3.2	
Choline, mg	80	
Biotin, mcg	8.0	
Pantothenic Acid, mg	1.3	
Inositol, mg	20	
Sodium, mg	90	
Potassium, mg	470	
Chloride, mg	230	
Calcium, mg	330	
Phosphorus, mg	250	



Magnesium, mg	40	
Iodine, mcg	23	
Manganese, mg	0.46	
Copper, mg	0.14	
Zinc, mg	1.7	
Iron, mg	2.7	
Selenium, mcg	8.0	
Chromium, mcg	9.0	
Molybdenum, mcg	9.0	



Instructions for Use:

8 fl oz Container

- Shake well.
- Store unopened at room temperature.
- Clean top of container thoroughly before opening.
- After opening, refrigerate and consume product within 48 hours. If consuming product directly from the container, refrigerate and consume within 24 hours.

Tube Feeding

- Follow physician's instructions for flow rate, volume, dilution and need for additional fluids.

Oral Feeding

- Best Served Chilled

Ready-To-Hang Container

All medical foods, regardless of type of administration system, require careful handling because they can support microbial growth. Follow these instructions for clean technique and proper setup to reduce the potential for microbial contamination.

NOTE: Failure to follow the **INSTRUCTIONS FOR USE** increases the potential for microbial contamination and may reduce hangtime.

- Administer product at room temperature.
- THOROUGHLY wash hands with soap and water before handling container or feeding set.
- Turn container upside down and SHAKE VIGOROUSLY, using a twisting motion for at least 10 seconds.
- DO NOT touch any part of the container or feeding set that comes into contact with the formula.
- When initiating feeding, follow physician's instructions. Adjust flow rate and volume according to patient's condition and tolerance.
- Additional fluid requirements should be met by giving water between or after feedings or when flushing the tube.

For Use with Enteral Feeding Pumps

- Remove dust cover from RTH Safety Screw Cap.
- Remove dust cover from feeding set connector.
- Insert feeding set connector into port of RTH Safety Screw Cap and completely pierce foil.
- Turn connector collar clockwise until it is securely fastened.
- Close clamp on set before inverting container.
- Invert container and suspend, using hanging ring on bottom of container.

Precautions

- Follow directions for use provided by manufacturer of feeding sets.
- Unless a shorter hang time is specified by the set manufacturer, hang product for up to 48 hours after initial connection when clean technique and only one new set are used. Otherwise hang for no more than 24 hours.