

Product Information: PediaSure® Enteral Formula 1.0 Cal With Fiber

For more information, contact your Abbott Nutrition Representative or visit
www.abbottnutrition.com

PediaSure® Enteral Formula 1.0 Cal With Fiber

Complete, Balanced Nutrition®



- 1 PEDIASURE ENTERAL FORMULA 1.0 CAL WITH FIBER is a source of complete, balanced nutrition especially designed for tube feeding children 1 to 13 years of age.
- 1 For tube feeding.
- 1 May be used as the sole source of nutrition or as a supplement.
- 1 Use under medical supervision.
- 1 No artificial growth hormones.*
- 1 Milk-based, complete, balanced nutrition.
- 1 1.0 Cal/mL, 240 Cal/8 fl oz, from a balanced distribution of protein, fat, and carbohydrate.
- 1 Meets or exceeds 100% of the DRIs for protein and 25 essential vitamins and minerals for children 1-8 years of age in 1000 Cal (1000 mL), and for children 9-13 years of age in 1500 Cal (1500 mL).
- 1 DHA (32 mg per 8 fl oz) and lutein to support the brain and eyes.^{1,2}
- 1 Blend of soluble and insoluble fibers and short-chain fructooligosaccharides at a level that is well tolerated by children.
- 1 Vitamin D, calcium and phosphorus for strong bones.
- 1 Prebiotics designed to support digestive tract health^{3,4,5,6} and antioxidants (vitamins C & E and selenium) to support the immune system.
- 1 Kosher.
- 1 Halal.
- 1 Gluten-free.
- 1 Suitable for lactose intolerance.

Safety Precautions

- 1 Not intended for infants under 1 year of age unless specified by a physician.
- 1 Not for IV use.
- 1 Not for children with galactosemia.

* No significant difference has been shown between milk derived from rbST-treated and non-rbST-treated cows.

¹ Lauritzen L, et al. *Prog Lipid Res.* 2001;40:1-94.
² Krinsky NI, et al. *Annu Rev Nutr.* 2003;23:171-201.
³ Bornet FR, et al. *Nutr Rev.* 2002;60:326-334.
⁴ Tokunaga T, et al. *Bifidus.* 1993;6:143-150.
⁵ Hidaka H, et al. *Bifidobact Microflora.* 1986;5:37-50.
⁶ Bouhnik Y, et al. *Nutr J* 2006; doi:10.1186/1 475-2891-5-8.

Ingredients

Liquid Vanilla:

Water, Corn Maltodextrin, Blend of Vegetable Oils (High Oleic Safflower, Soy, Medium Chain Triglycerides), Milk Protein Concentrate, Sugar, Soy Protein Isolate, Short-chain Fructooligosaccharides. Less than 0.5% of: Vitamins & Minerals (Potassium Phosphate, Calcium Carbonate, Potassium Chloride, Potassium Citrate, Calcium Phosphate, Ascorbic Acid, Choline Chloride, Ferrous Sulfate, dl-Alpha-Tocopheryl Acetate, Zinc Sulfate, Niacinamide, Calcium Pantothenate, Manganese Sulfate, Thiamine Hydrochloride, Pyridoxine Hydrochloride, Riboflavin, Vitamin A Palmitate, Copper Sulfate, Magnesium Phosphate, Folic Acid, Chromium Chloride, Potassium Iodide, Sodium Molybdate, Sodium Selenate, Biotin, Phylloquinone, Vitamin D3, Vitamin B12), Oat Fiber, Soy Fiber, Natural & Artificial Flavor, Gum Arabic, Cellulose Gel, Tuna Oil (Source of docosahexaenoic acid [DHA]), Cellulose Gum, Soy Lecithin, Monoglycerides, Carrageenan, Salt, Potassium Hydroxide, Inositol, Taurine, L-Carnitine, and Lutein.

Allergens: Contains milk and soy ingredients.

Availability

List Number	Item
67403	PediaSure Enteral 1.0 Cal with Fiber Vanilla / 8 fl oz (237 mL) Can / 24 ct
67405	PediaSure Enteral 1.0 Cal w Fiber Vanilla / 33.8 fl oz (1L) Ready-To-Hang Prefilled Container / 8 ct

For more information, contact your Abbott Nutrition Representative or visit www.abbottnutrition.com

PediaSure®

Enteral Formula 1.0 Cal With Fiber

Complete, Balanced Nutrition®

Nutrition Information - Liquid Vanilla

	8 fl oz (237 mL)		1000 mL (1 L)		1 L (Ped FC)		8 fl oz (Ped FC)	
	Value	%RDI / % DV	Value	%RDI / % DV	Value	%RDI / % DV	Value	%RDI / % DV
Volume, mL	237		1000		1000		237	
Protein, g	7		30		30		7	
Fat, g	9		38					
Saturated Fat, g	2.5		10		10		2.5	
Trans Fat, g	0		0		0		0	
Polyunsaturated Fat, g	2.5		11		11		2.5	
Monounsaturated Fat, g	3.5		15		15		3.5	
Cholesterol, mg	5		20		20		5	
Carbohydrate, g	34		143					
Dietary Fiber, g	3		13		13		3	
scFOS®, g	1.7		7		7		1.7	
Total Sugars, g	7		29		29		7	
Added Sugars, g	7		28		28		7	
L-Carnitine, mg	4		17		16.9		4	
Taurine, mg	18		76		76		18	
Water, g	200		845		845		200	
Calories	240		1000		1000		240	
Vitamin A, mcg RAE	140		590		590		140	
Vitamin D, mcg	6		25		25		6	
Vitamin E, mg	3		13		13		3	
Vitamin K, mcg	18		76		76		18	
Vitamin C, mg	23		97		97		23	
Folate, mcg DFE	60		253		253		60	
Folic Acid, mcg	36		152		152		36	
Thiamin (Vitamin B1), mg	0.3		1.3		1.3		0.3	
Riboflavin (Vitamin B2), mg	0.33		1.4		1.4		0.33	
Vitamin B6, mg	0.34		1.4		1.4		0.34	
Vitamin B12, mcg	0.48		2.0		2.0		0.48	
Niacin, mg NE	3.2		14		14		3.2	
Choline, mg	80		338		338		80	
Biotin, mcg	8		34		34		8	
Pantothenic Acid, mg	1.3		5.5		5.5		1.3	
Inositol, mg	20		84		84		20	
Sodium, mg	90		380					
Potassium, mg	470		1980					
Chloride, mg	230		970					
Calcium, mg	330		1390					
Phosphorus, mg	250		1050		1050		250	

For more information, contact your Abbott Nutrition Representative or visit www.abbottnutrition.com

PediaSure® Enteral Formula 1.0 Cal With Fiber

Complete, Balanced Nutrition®

Nutrition Information - Liquid Vanilla

	8 fl oz (237 mL)		1000 mL (1 L)		1 L (Ped FC)		8 fl oz (Ped FC)	
	Value	%RDI / % DV	Value	%RDI / % DV	Value	%RDI / % DV	Value	%RDI / % DV
Magnesium, mg	40		169		169		40	
Iodine, mcg	23		97		97		23	
Manganese, mg	0.46		1.9		1.9		0.46	
Copper, mg	0.14		0.59		0.59		0.14	
Zinc, mg	1.7		7.2		7.2		1.7	
Iron, mg	2.7		11		11		2.7	
Selenium, mcg	8		34		34		8	
Chromium, mcg	9		38		38		9	
Molybdenum, mcg	9		38		38		9	

Preparation

Instructions for Use:

8 fl oz can

- 1 Shake well.
- 1 Store unopened at room temperature.
- 1 Clean top of container thoroughly before opening.
- 1 After opening, cover, refrigerate and **use within 48 hours**.
- 1 Tube Feeding
 - 1 Follow physician's instructions for flow rate, volume, dilution and need for additional fluids.

Ready-To-Hang Container

- 1 All medical foods, regardless of type of administration system, require careful handling because they can support microbial growth. Follow these instructions for clean technique and proper setup to reduce the potential for microbial contamination. NOTE: Failure to follow the Instructions for Use increases the potential for microbial contamination and may reduce hangtime.
- 1 Administer product at room temperature.
- 1 THOROUGHLY wash hands with soap and water before handling container or feeding set.
- 1 Turn container upside down and SHAKE VIGOROUSLY, using a twisting motion for at least 10 seconds.
- 1 DO NOT touch any part of the container or feeding set that comes into contact with the formula.
- 1 When initiating feeding, follow physician's instructions. Adjust flow rate and volume according to patient's condition and tolerance.
- 1 Additional fluid requirements should be met by giving water between or after feedings or when flushing the tube.
- 1 Always follow directions for use provided with feeding set.
- 1 For Use with Enteral Feeding Set:
 - 1 Remove dust cover from RTH Safety Screw Cap.
 - 1 Remove dust cover from feeding set connector.
 - 1 Insert feeding set connector into port of Safety Screw Cap and completely pierce foil.
 - 1 Turn connector collar clockwise until it is securely fastened.
 - 1 Close clamp on set before inverting container.
 - 1 Invert container and suspend, using hanging ring on bottom of container.
 - 1 Unless a shorter hang time is specified by the set manufacturer, hang product for up to 48 hours after initial connection when clean technique and only one new set are used. Otherwise hang for no more than 24 hours.

For more information, contact your Abbott Nutrition Representative or visit www.abbottnutrition.com