

Drug Facts

DO NOT USE IF IMPRINTED SHRINKBAND IS MISSING OR BROKEN.

Active ingredient (in each 30 mL) Purpose

Bismuth subsalicylate 1050 mgUpset stomach reliever and anti-diarrheal

Uses relieves: ■ diarrhea ■ heartburn ■ indigestion ■ nausea ■ upset stomach associated with these symptoms

Warnings

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Contains salicylate. Do not take if you are

- allergic to salicylates (including aspirin)
- taking other salicylate products

Do not use if you have ■ an ulcer

- bloody or black stool ■ a bleeding problem

Ask a doctor before use if you have

- fever ■ mucus in the stool

Ask a doctor or pharmacist before use if you are

- taking any drug for ■ anticoagulation (thinning of the blood) ■ diabetes ■ gout ■ arthritis

When using this product a temporary, but harmless darkening of the stool and/or tongue may occur

Stop use and ask a doctor if ■ symptoms get worse

- ringing in the ears or loss of hearing occurs
- diarrhea lasts more than 2 days

If pregnant or breast-feeding, ask a health professional before use. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center immediately.

◀ **Drug Facts continued**

MAJOR®

Maximum Strength Bismatrol

**Soothing, coating action provides
fast relief for heartburn,
indigestion, nausea,
upset stomach and diarrhea**

**COMPARE TO THE ACTIVE
INGREDIENT IN PEPTO-BISMOL®.***

**8 FL OZ
(236 mL)**

NDC 0904-1314-09

Drug Facts (continued)

Directions

 ■ shake well before use

■ mL = milliliter ■ TBSP = tablespoon ■ adults and children 12 years and over: 1 dose (2 TBSP or 30 mL) every hour as needed ■ do not exceed 4 doses (8 TBSP or 120 mL) in 24 hours ■ use until diarrhea stops but not more than 2 days ■ children under 12 years: ask a doctor ■ drink plenty of clear fluids to help prevent dehydration caused by diarrhea

Other information

 ■ each 30 mL or 2 TBSP contains:

- potassium 25 mg • salicylate 471 mg • sodium 8 mg
- protect from freezing. ■ avoid excessive heat (over 104°F or 40°C). ■ dosage cup provided

Inactive ingredients benzoic acid, D&C red # 22, D&C red # 28, flavor, hydroxyethyl cellulose, potassium hydroxide, purified water, saccharin sodium, salicylic acid, simethicone, xanthan gum.

Questions or Comments? (800) 616-2471

*This product is not manufactured or distributed by Procter & Gamble, Inc., the distributor of Pepto-Bismol® Maximum Strength.

Distributed By MAJOR® PHARMACEUTICALS

17177 N. Laurel Park Drive, Suite 233
Livonia, MI 48152 USA

Re-Order No. 700283
M-30



MFR# 53041 REV 01/18