

Mintox Plus Tabs

Dosage form: tablet, chewable

Ingredients: ALUMINUM HYDROXIDE 200mg, MAGNESIUM HYDROXIDE 200mg, SIMETHICONE 25mg

Labeler: Major Pharmaceuticals

NDC Code: 0904-0478

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

Mintox Plus Tabs-Antacid Plus Anti-Gas

Active ingredients (in each tablet)

Aluminum hydroxide (equiv. to dried gel, USP) 200 mg

Magnesium hydroxide 200 mg

Simethicone 25 mg

Purpose

Antacid

Antacid

Antigas

Uses

relieves:

- acid indigestion
- heartburn
- sour stomach
- upset stomach and gas associated with these symptoms

Warnings

Ask a doctor before use if you have

- kidney disease
- a magnesium-restricted diet

Ask a doctor or pharmacist before use if you are

- presently taking a prescription drug. Antacids may interact with certain prescription drugs.

Do not take more than 16 tablets in a 24-hour period, or use the maximum dosage of this product for more than 2 weeks, except under the advice and supervision of a doctor.

Keep out of reach of children. In case of overdose get medical help or contact a poison control center immediately.

Directions

- chew 1 to 4 tablets 4 times a day or as directed by a doctor.

Other information

- each tablet contains: magnesium 85 mg
- do not use if printed seal under cap is torn or missing
- store at room temperature
- keep tightly closed and dry

Inactive ingredients

citric acid anhydrous, corn starch, D and C red 30 lake, D and C yellow 10 lake, dextrose, flavors, glycerin, magnesium stearate, maltodextrin, mannitol, pregelatinized starch, saccharin sodium, sorbitol, sugar

Principal Display Panel

MAJOR[®]

Mintox Plus Tabs

Antacid plus Anti-gas

Fast Relief of:

- Acid Indigestion
- Heartburn
- Sour stomach
- Gas

Great Taste!

Compare to the Active ingredients in Maalox[®] Plus
100 Tablets

NDC 0904-0478-60



MintoxTM Plus Tabs

ANTACID PLUS ANTI-GAS

FAST RELIEF OF:

- ACID INDIGESTION
- HEARTBURN
- SOUR STOMACH
- GAS



GREAT TASTE!

100 TABLETS

**Compare to the Active Ingredients in Maalox[®] Plus*

Drug Facts

Active ingredients (in each tablet)

| Active ingredients (in each tablet) | Purpose |
|------------------------------------------------------|---------|
| Aluminum hydroxide (equiv. to dried gel, USP) 200 mg | Antacid |
| Magnesium hydroxide 200 mg | Antacid |
| Simethicone 25 mg | Antigas |

Uses relieves: ■ acid indigestion ■ heartburn ■ sour stomach ■ upset stomach & gas associated with these symptoms

Warnings

Ask a doctor before use if you have ■ kidney disease ■ a magnesium-restricted diet
 Ask a doctor or pharmacist before use if you are ■ presently taking a prescription drug. Antacids may interact with certain prescription drugs.

Do not take more than 16 tablets in a 24-hour period, or use the maximum dosage of this product for more than 2 weeks, except under the advice and supervision of a doctor.

Keep out of reach of children. In case of overdose get medical help or contact a Poison Control Center immediately.

Directions ■ chew 1 to 4 tablets 4 times a day or as directed by a doctor

Other information ■ each tablet contains: magnesium 85 mg ■ do not use if printed seal under cap is torn or missing ■ store at room temperature ■ keep tightly closed and dry

Inactive ingredients citric acid anhydrous, corn starch, D&C red 30 lake, D&C yellow 10 lake, dextrose, flavors, glycerin, magnesium stearate, maltodextrin, mannitol, pregelatinized starch, saccharin sodium, sorbitol, sugar.

Questions or Comments? (800) 616-2471

*This product is not manufactured or distributed by Novartis Consumer Health Inc., owner of the registered trademark MAALOX[®] Plus.

Distributed By: MAJOR[®] PHARMACEUTICALS
 31778 Enterprise Drive, Livonia, MI 48150 USA

LOT: M-30
 M-30
 Re-Order No. 700803
 MFN# 53041 REV 04/12



EXP:

MINTOX PLUS TABS

aluminum hydroxide, magnesium hydroxide, simethicone tablet, chewable

Product Information

| | | | |
|-------------------------|----------------|--------------------|---------------|
| Product Type | HUMAN OTC DRUG | Item Code (Source) | NDC:0904-0478 |
| Route of Administration | ORAL | DEA Schedule | |

Active Ingredient/Active Moiety

| Ingredient Name | Basis of Strength | Strength |
|----------------------------------------------------------|---------------------|----------|
| ALUMINUM HYDROXIDE (ALUMINUM HYDROXIDE) | ALUMINUM HYDROXIDE | 200 mg |
| MAGNESIUM HYDROXIDE (MAGNESIUM CATION and HYDROXIDE ION) | MAGNESIUM HYDROXIDE | 200 mg |
| SIMETHICONE (SIMETHICONE) | SIMETHICONE | 25 mg |

Inactive Ingredients

| Ingredient Name | Strength |
|----------------------------|----------|
| ANHYDROUS CITRIC ACID | |
| STARCH, CORN | |
| D&C RED NO. 30 | |
| D&C YELLOW NO. 10 | |
| DEXTROSE | |
| GLYCERIN | |
| MAGNESIUM STEARATE | |
| MALTODEXTRIN | |
| MANNITOL | |
| SACCHARIN SODIUM DIHYDRATE | |

SORBITOL

SUCROSE

Product Characteristics

| | | | |
|-------|--------|-------|----------|
| Color | yellow | Score | no score |
|-------|--------|-------|----------|

| | | | |
|-------|-------|------|------|
| Shape | ROUND | Size | 16mm |
|-------|-------|------|------|

| | | | |
|--------|-------|--------------|------|
| Flavor | LEMON | Imprint Code | G152 |
|--------|-------|--------------|------|

Contains

Packaging

| # | Item Code | Package Description |
|---|------------------|-------------------------------------------|
| 1 | NDC:0904-0478-60 | 100 TABLET, CHEWABLE (TABLET) in 1 BOTTLE |

Marketing Information

| Marketing Category | Application Number or Monograph Citation | Marketing Start Date | Marketing End Date |
|---------------------|------------------------------------------|----------------------|--------------------|
| OTC monograph final | part331 | 03/14/2007 | |

Labeler - Major Pharmaceuticals (191427277)

Revised: 08/2012
Major Pharmaceuticals

Disclaimer: Every effort has been made to ensure that the information provided here is accurate, up-to-date and complete, but no guarantee is made to that effect. Drug information contained herein may be time sensitive. This information has been compiled for use by healthcare practitioners and consumers in the United States. The absence of a warning for a given drug or combination thereof in no way should be construed to indicate that the drug or combination is safe, effective or appropriate for any given patient. If you have questions about the substances you are taking, check with your doctor, nurse or pharmacist.