



UTI-Stat®

A ready-to-drink medical food providing cranberry concentrate with added ingredients (vitamin C, D-Mannose, fructo-oligosaccharides (FOS), and bromelain) in a 1 fl oz serving for urinary tract health.

UTI-Stat contains 2 % cranberry juice concentrate.

Main Features:

- Clinically supported in 2 published trials to manage urinary tract infections^{1,2}
- Low in sugar (1 g per serving)
- Lactose-, gluten-, and soy-free
- Kosher
- Available in cranberry flavor
- For oral and enteral use

Indications:

- Dietary management of urinary tract infections
- Urinary tract health

Ordering/Reimbursement Information:

Product Code	Reimbursement Code	HCPCS Code	Product Packaging	Calories Per Unit
78357 - Cranberry (bottle)	26974-0410-66	B4104	4 x 30 fl oz	2100
78404 - Cranberry (unit)	26974-0410-67	B4104	96 x 1 fl oz	70

Nutrition Information:

Nutrients	Per 30 mL (2 Tbsp)
Serving Size: 1 fl oz	
Calories	70*
Sugar, g	1
Minerals	
Sodium, mg	15
Potassium, mg	20
Phosphorus, mg	0
<i>Proprietary Blend (Cranberry Concentrate, Ascorbic Acid, D-Mannose, FOS, Bromelain)</i>	
<i>*Glycerin provides a source of calories</i>	

Ingredients:

Water, Glycerin, Proprietary Blend (Cranberry Concentrate, Ascorbic Acid, D-Mannose, Fructo-oligosaccharides (FOS), Bromelain), Contains 2 % or less: Sucralose, Acesulfame Potassium, Potassium Sorbate (to preserve flavor), Sodium Benzoate (to preserve flavor), Phosphoric Acid.

Administration:

- Use under medical supervision.
- Administer orally or through feeding tube.

Suggested Intake:

- 1-2 servings per day as needed.
- One serving = 1 fl oz.

Oral Feeding:

- Can be taken straight or mixed with any hot or cold food or beverage.

Tube Feeding:

1. Flush feeding tube with 30-60 mL water.
2. Measure and administer dose prescribed.
3. Administer UTI-Stat straight into the enteral tube.
4. Flush the tube with additional 30 mL water.

Precautions:

- Not a substitute for antibiotics.
- This product is not intended for use as an analgesic.
- Not suitable for use as a sole source of nutrition.
- Not for parenteral use.

Storage:

- Store at room temperature.
- Refrigeration not necessary.
- Discard 3 months after opening.
- Record date opened on bottom of container.
- Do not use if seal is broken.

¹ Pothauer ME, The Director. 2008;16:13-6. ² Efron M, et al. Urology. 2010;75:841-5.