



Pro-Stat® Renal Care

A sugar free, ready-to-drink liquid protein medical food providing 15 g of enzyme-hydrolyzed complete protein, 3 g of prebiotic fiber and 100 calories per 1 fl oz.

Main Features:

- Clinically supported in 1 published trial to improve protein anabolism in hemodialysis patients¹
- 15 g of complete protein and 3 g of prebiotic soluble fiber in 1 fl oz
- Hydrolyzed for rapid absorption and efficient utilization²
- High in conditionally-indispensable amino acids needed for recovery³⁻⁸
- Low volume making it appropriate for fluid restricted patients
- Sugar free
- Honey-like viscosity
- Lactose free, gluten free, and soy free
- Kosher
- Available in great tasting tangerine flavor
- Available in bottles (6/case) or unit doses (96/case)
- For oral and enteral use

Indications:

- Increased protein needs in low volume related to:
 - Pressure injuries
 - Wounds (diabetic, venous, surgical, burns)
- Dialysis
- Fluid restriction
- Unintentional muscle loss
- Protein-energy malnutrition
- Low serum proteins
- Sarcopenia
- Inadequate protein & fiber intake

Ordering/Reimbursement Information:

Product Code	Reimbursement Code	HCPCS Code	Product Packaging	Calories Per Unit
78388 - Tangerine (bottle)	26974-0410-64	B4155	6 x 30 fl oz	3000
78406 - Tangerine (unit)	26974-0420-65	B4155	96 x 1 fl oz	100

Nutrition Information:

Nutrients	Per 30 mL (2 Tbsp)
Calories	100
Protein, g	15
Fat, g	0
Carbohydrate, g	6
Fiber, g	3
Minerals	
Sodium, mg	50
Potassium, mg	20
Phosphorus, mg	50

Ingredients:

Hydrolyzed Collagen Protein, Water, Glycerin, Polydextrose, Fructo-Oligosaccharides (FOS), Malic Acid. Contains 2% or Less: Natural & Artificial Flavor, Phosphoric Acid, L-Tryptophan, Sucralose, Potassium Sorbate (to help protect flavor), Sodium Benzoate (to help protect flavor).

Administration:

- Use under medical supervision.
- Healthcare professionals should determine intake based on condition, estimated need, and dietary intake.
- Administer orally or through feeding tube.

Suggested Intake:

- 1-3 servings per day.
- One serving = 1 fl oz.

Oral Feeding:

- Shake well before opening.
- Can be taken straight or mixed with any hot or cold food or beverage.

Tube Feeding:

1. Flush feeding tube with 30-60 mL of water.
2. Pour 30 mL of Pro-Stat in a 4-6 fl oz container.
3. Add 30-60 mL water and mix well.
4. Administer Pro-Stat via syringe.
5. Flush with 30-60 mL of water.

Precautions:

- Do not add Pro-Stat to an open or closed system of the tube feeding formula or into a container of enteral feeding.
- Do not add medications to Pro-Stat, even after dilution.
- Not suitable for use as a sole source of nutrition.
- Not for parenteral use.

Storage:

- Store at room temperature.
- Refrigeration causes gelling.
- Discard 3 months after opening.
- Record date opened on bottom of container.
- Do not use if seal is broken.

1. Sundell MB, et al. J Ren Nutr. 2009;19:412-21. 2. Koopman R, et al. Am J Clin Nutr. 2009;90:106-15. 3. Morris Jr, SM. Annu Rev Nutr. 1992;12:81-101. 4. Curis E, et al. Amino Acids. 2005;29:177-205. 5. Wu G, et al. Biochem J. 1992;281:45-8. 6. Schwedhelm E, et al. Br J Clin Pharmacol. 2008;65:51-9. 7. Waugh WH, et al. Am J Hematol. 1999;61:216. 8. Waugh WH, et al. J Natl Med Assoc. 2001;93:363-71.