

GERI CARE®

ADULT LOW DOSE
CHEWABLE**ASPIRIN 81 mg**

PAIN RELIEVER (NSAID)

Compare to active ingredient in
BAYER® Low Dose Aspirin®

36 Tablets

Tamper Evident: Do not use if imprinted seal under cap is missing or broken.**Drug Facts**

Purpose
Active ingredient (in each tablet) Aspirin 81 mg (NSAID)* Pain reliever
 *nonsteroidal anti-inflammatory drug

Uses for the temporary relief of minor aches and pains or as recommended by your doctor

Warnings

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Aspirin may cause a severe allergic reaction which may include: ■ hives ■ facial swelling

■ shock ■ asthma (wheezing)

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you: ■ are age 60 or older ■ have had stomach ulcers or bleeding problems ■ take a blood thinning (anticoagulant) or steroid drug ■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) ■ have 3 or more alcoholic drinks every day while using this product ■ take more or for a longer time than directed

Do not use ■ if you are allergic to aspirin or any other pain reliever/fever reducer ■ this product for at least 7 days after tonsillectomy or oral surgery unless directed by a doctor ■ if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if ■ stomach bleeding warning applies to you ■ you have a history of stomach problems, such as heartburn ■ you have high blood pressure, heart disease, liver cirrhosis, or kidney disease ■ you are taking a diuretic ■ you have asthma

* This product is not manufactured or distributed by the owner of the registered trademark Bayer®. Dist. by: **Geri-Care Pharmaceuticals Corp.**
 1650 63rd Street, Brooklyn, NY 11204

REV 0721



3 57896 91136 2

PEEL HERE FOR MORE DRUG FACTS →

STOP PEELING

ADHESIVE

Questions or comments? 1-800-540-3765

saccharin sodium.

FD&C yellow #6 aluminum lake, orange flavor,

Inactive ingredients: corn starch, dextarates,

15°C - 30°C (59°F - 86°F)

Other information ■ store at room temperature

ask a doctor

children under 12

or as directed by a doctor

than 48 tablets in 24 hours,

4 hours as needed, not more

12 years and older

Take 4-8 tablets every

do not exceed recommended dose

■ do not drink alcohol while taking

■ drink a full glass of water with each dose

Keep out of reach of children. In case of

overdose, get medical help or contact a Poison

Control Center (1-800-222-1222) right away.

Keep out of reach of children. In case of

overdose, get medical help or contact a Poison

Control Center (1-800-222-1222) right away.

Keep out of reach of children. In case of

overdose, get medical help or contact a Poison

Control Center (1-800-222-1222) right away.

Keep out of reach of children. In case of

overdose, get medical help or contact a Poison

Control Center (1-800-222-1222) right away.

Keep out of reach of children. In case of

overdose, get medical help or contact a Poison

Control Center (1-800-222-1222) right away.

Keep out of reach of children. In case of

overdose, get medical help or contact a Poison

Control Center (1-800-222-1222) right away.

Keep out of reach of children. In case of

overdose, get medical help or contact a Poison

Control Center (1-800-222-1222) right away.

Keep out of reach of children. In case of

overdose, get medical help or contact a Poison

Control Center (1-800-222-1222) right away.

Drug Facts (continued)

Ask a doctor or pharmacist before use if you are

taking a prescription drug for

■ gout ■ diabetes ■ arthritis

Stop use and ask a doctor if ■ an allergic

reaction occurs. Seek medical help right away

■ you experience any of the following signs of

stomach bleeding ■ feel faint ■ vomit blood

■ have bloody or black stool

■ pain gets worse or lasts more than 10 days

■ redness or swelling is present

■ new symptoms occur

■ ringing in the ears or loss of hearing occurs